## Hydrofera Blue Classic

Polyvinyl Alcohol (PVA) Foam Dressings

## Hydrofera Blue READY and READY-Transfer

Polyurethane (PU) Foam Dressings

## Hydrofera Blue READY-Border

Polyurethane Foam Dressing with Silicone Border

#### **Wound Characteristics**

- Deep, tunneling or shallow
- With or without devitalized tissue/ slough & debris
- Moderate to heavy exudate
- Suspected biofilm
- Infected or at risk of infection
- Safe, Non-cytotoxic Does not inhibit growth factors

• Safe, autolytic debridement

Continuous wicking, highly absorptive

High tensile strength

Manages bioburden

Clinical Benefits

Kills bacteria in the dressing



### **Wound Characteristics** Shallow wounds Moderate to heavy exudate Clinical Benefits

- Manages bioburden Safe, Non-cytotoxic
- Kills bacteria in the dressing

### • Up to 7-day wear time



#### Wound Characteristics:

- Shallow wounds
- Moderate to heavy exudate
- Infected wounds or wounds at risk of infection


### Clinical Benefits:

- Non-cytotoxic
- Does not inhibit growth factors Gentle, secure hold
- Atraumatic removal
- Optimal moist wound healing
- Up to 7-day wear time

• Kills bacteria in the dressing

#### **Powerful Wicking**

• Wicks bacteria laden exudate into the foam and away from the wound surface with natural negative pressure through capillary flow.



### Edge Effect

• Multiple authors have noted resolution of epibole or flattening of wound edges with use of Hydrofera Blue CLASSIC dressings. 1-4







Granulating wound bed with epithelial cells on the edge

#### Debridement

• Devitalized tissue removed at dressing change.




#### The Safe Non-Cytotoxic Choice

- Ideal for clinicians looking for an alternative to silver dressings
- Less expensive than many silver foam dressings<sup>5</sup>
- Compatible for use with Hyperbaric Oxygen Therapy (HBOT)
- Compatible with Growth factors and enzymatic debriders





Absorbs and retains bacterialaden exudate





### **READY-Transfer** dressing No-film backing increases exudate management options Exudate is transferred into secondary absorbent dressing



Secure Fit for difficult to dress areas and mobility



Diabetic foot ulcers



yet gentle removal

Intimate contact of foam to the wound bed










Deep with Heavy Exudate



With Disposable NPWT



Removal of Devitalized Tissue



**Shallow Wounds** 



Skin Tears



**Under Total Contact Casting** 



With Compression



Leg Ulcer



Atraumatic removal


## Hydrofera Blue Specialty Dressings

### **Tunneling Dressing**



- A unique antimicrobial wound dressing that can be used for Sinuses, Tunnels and Negative Pressure Wound Therapy
- Wicks bacteria laden execute, slough and debris out of the wound
- High tensile strength, easy one-piece removal
- Wet the tip prior to insertion

### Ostomy Dressing



- Manages bioburden, which helps the bodies own immune system restore bacterial balance
- Broad spectrum activity against a wide variety of bacteria and yeast commonly found in wounds
- No hydration required

### Hydrofera Blue helps wounds heal faster at less cost<sup>6</sup>

A recent retrospective study examining wound treatment data of more than 30,000 patients over a two year period, Hydrofera Blue dressings combined with an Advanced Wound Management Program showed faster healing times and lower treatment costs.

- Compared with other advanced wound dressings, Hydrofera Blue showed a 7-day reduction in healing time and 24% reduction in nursing Labour cost.
- Compared with other commonly used dressings on patients who were not on an Advanced Wound Management Program, Hydrofera Blue delivered over a 50% reduction in healing time and a 75% reduction in nursing labour cost.

#### All Hydrofera Blue dressings are effective against a broad-spectrum of microorganisms commonly found in wounds.

Bacteria				Yeast
Methicillin-resistant Staphylococcus aureus (MRSA) Staphylococcus aureus	Pseudomonas fluorescens	Klebsiella pneumoniae	Bacillus subtilis	Candida albicans
			Yersinia enterocolitica	Candida krusei
	Escherichia coli		Serratia marcescens	Candida glabrata
	Enterococcus faecalis	Proteus mirabilis		O .
Staphylococcus epidermidis (coagulase negative)	Vancomycin-resistant Enterococcus (VRE)	Proteus vulgaris	Streptococcus pneumoniae**	
Pseudomonas aeruginosa	Enterococcus (VRE)	Enterobacter aerogenes	Enterobacter cloacae**	

<sup>\*</sup> Tested on Hydrofera Blue Ready dressing only

### **Ordering Information**

	Hydrofera Blue <b>CL</b>	ASSIC® Antibacterial Dressings		
	Product No	Product Size	Qty/Box	HCPCS*
	Standard Dressings HB2214 HB4414 HB4514 HB6614 HBT0906	2" x 2" (5 cm x 5 cm) 4" x 4" (10.2 cm x 10.2 cm) 4" x 5" (10.2 cm x 12.7 cm) 6" x 6" (15.2 cm x 15.2 cm) 9 mm Tunneling Dressing (1.2 g)	10 10 10 10 10	A6209 A6209 A6210 A6210 A6215
	Heavy Drainage Dres HBHD4450 HBHD6675	ssings 4" x 4" x 0.5" thick (10.2 cm x 10.2 cm x 1.2 cm) 6" x 6" x 0.75" (15.2 cm x 15.2 cm x 1.9 cm)	5 5	A6209 A6210
	Dressings with Moist HBRF4414 HBRF2258 HBRF4475	ture-Retentive Film  4" x 4" (10.2 cm x 10.2 cm)  2.25" x 8" (5.7 cm x 20.3 cm)  4" x 4.75" (10.2 cm x 12 cm) Island Dressing;  pad size 2" x 2.75" (5 cm x 6.5 cm)  2.5" (6cm) Diameter Ostomy Dressing	10 10 10	A6209 A6210 A6212 A6209
	Hvdrofera Blue <b>RE</b>	ADY® Antibacterial Dressings		
SYEADY 19 19  Thydroless Date Focador 19  There District 19 19 19 19 19 19 19 19 19 19 19 19 19	HBRS2520 HBRS4520 HBRS8820	2.5" x 2.5" (6.4 cm x 6.4 cm) 4" x 5" (10.2 cm x 12.7 cm) 8" x 8" (20.3 cm x 20.3 cm)	10 10 10	A6209 A6210 A6211
	Hydrofera Blue RI	EADY®-Transfer Antibacterial Dressings		
	HBRT2525 HBRT4050 HBRT8080	2.5" x 2.5" (6.4 cm x 6.4 cm) 4" x 5" (10.2 cm x 12.7 cm) 8" x 8" (20.3 cm x 20.3 cm)	10 10 10	A6209 A6210 A6211
	Hydrofera Blue RI	EADY®-Border Dressings		
NEW  SECTION OF THE PROPERTY O	HBRB4040 HBRB5959 HBRB6080	4" x 4" (10.2 cm x 10.2 cm) 5.9" x 5.9" (15 cm x 15 cm) 6" x 8" (15.2 cm x 20.3 cm)	10 10 10	A6212 A6212 A6213
READY TO HYDR	*1.100.00			

\*HCPCS are for reference only and subject to change.

Hydrofera Blue Dressings are represented by Appulse


Change to blue at hydrofera.com

Indications for use: Pressure Ulcers, Diabetic Ulcers, Venous Statis Ulcers, Arterial Ulcers, Superficial Burns Donor Sites, Skin Grafts, Post Surgical Incisions, Traumatic Wounds, Skin Tears, Abrasions, Lacerations,

- 3. Weir D, Blakely M. Case review of the clinical use of an antimicrobial PVA foam dressing. Poster presented at Symposium on Advances in Skin and Wound Care; April 18-21, 2012; Atlanta, GA.
- 4. Convell P, Mikulski L, Tramontozzi M. A comparison of two antimicrobial PVA foam dressings: a randomized prospective trial comparing PVA foam with two organic pigments to a silver wound dressing. Poster presented at Symposium on Advanced Wound Care, May 2-5, 2004; Lake Buena Vista, Fla.
- 5. Hurd T. Improving the quality of chronic wound care using an advanced wound management program and Gentian Violet / Methylene Blue-Impregnated Antibacterial Dressings: A Retrospective Study. Surgical Technology International Volume 35, Sept 2019

#### For product questions and samples please call Customer Service at 1-860-337-7730.

Hydrofera Dressings manufactured by: www.hvdrofera.com



# The most trusted line of foam dressings gives rise to a new bue



**NEW** Hydrofera Blue READY®-Border provides a perfect solution for secure wound protection and atraumatic dressing removal.





